

Policy # 00023 Original Effective Date: 01/26/2004 Current Effective Date: 08/12/2024

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

Note: Whole Gland Cryoablation of Prostate Cancer is addressed separately in medical policy 00022.

Note: Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors is addressed separately in medical policy 00175.

Note: Radiofrequency Ablation of Primary or Metastatic Liver Tumors is addressed separately in medical policy 00182.

Note: Cryosurgical Ablation of Primary or Metastatic Liver Tumors is addressed separately in medical policy 00220.

When Services Are Eligible for Coverage

Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:

- Benefits are available in the member's contract/certificate, and
- Medical necessity criteria and guidelines are met.

Based on review of available data, the Company may consider cryosurgical ablation for clinically localized, suspected renal malignancy for individuals with peripheral lesions that are less than or equal to 4 cm in diameter to be **eligible for coverage.****

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisianais an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00023 Original Effective Date: 01/26/2004 Current Effective Date: 08/12/2024

When Services May Be Eligible for Coverage

Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:

- Benefits are available in the member's contract/certificate, and
- Medical necessity criteria and guidelines are met.

Based on review of available data, the Company may consider cryosurgical ablation to treat lung cancer to be **eligible for coverage.****

Patient Selection Criteria

Coverage eligibility for cryosurgical ablation to treat lung cancer may be considered when **EITHER** of the following criteria is met:

- The individual has early-stage non-small cell lung cancer (NSCLC) and is a poor surgical candidate; **OR**
- The individual requires palliation for a central airway obstructing lesion.

When Services Are Considered Investigational

Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.

Based on review of available data, the Company considers cryosurgical ablation to treat localized renal cell carcinoma (RCC) in all other situations to be **investigational.***

Based on review of available data, the Company considers cryosurgical ablation to treat lung cancer when patient selection criteria are not met to be **investigational.***

Based on review of available data, the Company considers cryosurgical ablation as a treatment for benign or malignant tumors of the breast, pancreas, or bone to be **investigational.***

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisianais an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00023 Original Effective Date: 01/26/2004 Current Effective Date: 08/12/2024

Policy Guidelines

The objective of this review is to determine whether cryoablation of tumors located in the kidney, lung, breast, pancreas, or bone will improve the net health outcome. This review is limited to treatment in adults (age 18 years and older) and does not address pediatric populations.

Background/Overview

Renal Tumors

Localized kidney cancer is treated with radical nephrectomy or nephron-sparing surgery. Prognosis drops precipitously if the tumor extends outside the kidney capsule because chemotherapy is relatively ineffective against metastatic renal cell carcinoma.

Lung Tumors and Lung Metastases

Early-stage lung tumors are typically treated surgically. Patients with early-stage lung cancer who are not surgical candidates may be candidates for radiotherapy with curative intent. Cryoablation is being investigated in patients who are medically inoperable, with small primary lung cancers or lung metastases from extrapulmonary primaries. Patients with a more advanced local disease or metastatic disease may undergo chemotherapy with radiation following resection. Treatment is rarely curative; rather, it seeks to retard tumor growth or palliate symptoms.

Breast Tumors

Early-stage primary breast cancers are treated surgically. The selection of lumpectomy, modified radical mastectomy, or another approach is balanced against the patient's desire for breast conservation, the need for tumor-free margins in resected tissue, and the patient's age, hormone receptor status, and other factors. Adjuvant radiotherapy decreases local recurrences, particularly for those who select lumpectomy. Adjuvant hormonal therapy and/or chemotherapy are added, depending on the presence and number of involved nodes, hormone receptor status, and other factors. Treatment of metastatic disease includes surgery to remove the lesion and combination chemotherapy.

Fibroadenomas are common benign tumors of the breast that can present as a palpable mass or a mammographic abnormality. These benign tumors are frequently surgically excised to rule out a malignancy.

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisianais an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00023 Original Effective Date: 01/26/2004 Current Effective Date: 08/12/2024

Pancreatic Cancer

Pancreatic cancer is a relatively rare solid tumor that occurs almost exclusively in adults, and it is largely considered incurable. Surgical resection of tumors contained entirely within the pancreas is currently the only potentially curative treatment. However, the nature of the cancer is such that few tumors are found at such an early and potentially curable stage. Patients with a more advanced local disease or metastatic disease may undergo chemotherapy with radiation following resection. Treatment focuses on slowing tumor growth and palliation of symptoms.

Bone Cancer and Bone Metastases

Primary bone cancers are extremely rare, accounting for less than 0.2% of all cancers. Bone metastases are more common, with clinical complications including debilitating bone pain. Treatment for bone metastases is performed to relieve local bone pain, provide stabilization, and prevent impending fracture or spinal cord compression.

FDA or Other Governmental Regulatory Approval

U.S. Food and Drug Administration (FDA)

Several cryoablation devices have been cleared for marketing by the U.S. Food and Drug Administration (FDA) through the 510(k) process for use in open, minimally invasive, or endoscopic surgical procedures in the areas of general surgery, urology, gynecology, oncology, neurology, dermatology, proctology, thoracic surgery, and ear, nose, and throat. Examples include:

- Cryocare^{®‡} Surgical System (Endocare);
- CryoGen Cryosurgical System (Cryosurgical);
- CryoHit^{®‡} (Galil Medical) for the treatment of breast fibroadenoma;
- IceSense3^{™‡}, ProSense^{™‡}, and MultiSense Systems (IceCure Medical);
- SeedNet[™][‡] System (Galil Medical); and
- Visica^{®‡} System (Sanarus Medical).

FDA product code: GEH.

Rationale/Source

This medical policy was developed through consideration of peer-reviewed medical literature generally recognized by the relevant medical community, U.S. Food and Drug Administration

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisianais an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00023 Original Effective Date: 01/26/2004 Current Effective Date: 08/12/2024

approval status, nationally accepted standards of medical practice and accepted standards of medical practice in this community, technology evaluation centers, reference to federal regulations, other plan medical policies, and accredited national guidelines.

Description

Cryosurgical ablation (hereafter referred to as cryosurgery or cryoablation) involves freezing of target tissues; this is most often performed by inserting a coolant-carrying probe into the tumor. Cryosurgery may be performed as an open surgical technique or as a closed procedure under laparoscopic or ultrasound guidance.

Summary of Evidence

For individuals with early stage kidney cancer who are surgical candidates treated with cryoablation, the evidence includes comparative observational studies and systematic reviews. Relevant outcomes are overall survival (OS), disease-specific survival, quality of life, and treatment-related morbidity. Multiple comparative observational studies and systematic reviews of these studies have compared cryoablation to partial nephrectomy for early stage renal cancer. These studies have consistently found that partial nephrectomy is associated with better oncological outcomes than cryosurgery, but cryosurgery was associated with better perioperative outcomes, lower incidence of complications, and less decline in kidney function. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with early stage kidney cancer who are not surgical candidates and who are treated with cryoablation, the evidence includes comparative observational studies of cryoablation compared to partial nephrectomy or other ablative techniques, systematic reviews of these studies, and case series. Relevant outcomes are OS, disease-specific survival, quality of life, and treatment-related morbidity. Although oncological outcomes were better with surgery, in comparative observational studies, cryoablation was associated with less decline in kidney function. Recent case series totaling more than 400 patients showed cryoablation was associated with good oncological outcomes and preservation of renal function. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with non-small cell lung cancer (NSCLC) who are not surgical candidates, the evidence includes uncontrolled observational studies and case series. Relevant outcomes are OS, disease-specific survival, quality of life, and treatment-related morbidity. Medically inoperable

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisianais an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Policy # 00023 Original Effective Date: 01/26/2004 Current Effective Date: 08/12/2024

patients with early stage primary lung tumors were treated with cryoablation in a consecutive series of 45 patients. Five year survival was 68%; the main complications were hemoptysis in 40% of patients and pneumothorax in 51%. A prospective single arm Phase 2 study of 128 patients reported on cryoablation for treatment of metastases to the lung. Cryoablation for metastatic lung cancer was studied in a single arm trial in 40 patients. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with NSCLC who require palliation for a central airway obstructing lesion who are treated with cryoablation, the evidence includes case series. Relevant outcomes are OS, disease-specific survival, quality of life, and treatment-related morbidity. There are no comparative studies. A series of 521 consecutive patients reported improvement in symptoms in 86% of patients, but multiple study design, conduct, and relevance limitations preclude drawing conclusions about efficacy or safety of cryoablation in this population. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with solid tumors located in the breast, pancreas, or bone who are treated with cryoablation, the evidence includes uncontrolled observational studies and case series. Relevant outcomes are OS, disease-specific survival, quality of life, and treatment-related morbidity. Due to the lack of prospective controlled trials, it is not possible to conclude that cryoablation improves outcomes for any indication better than alternative treatments. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Supplemental Information

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

Clinical Input From Physician Specialty Societies and Academic Medical Centers

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

^{©2024} Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisianais an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Policy # 00023 Original Effective Date: 01/26/2004 Current Effective Date: 08/12/2024

2017 Input

Clinical input was sought to help determine whether the use of cryoablation for individuals with non small cell lung cancer (NSCLC) who are either poor surgical candidates or who required palliation for a lesion obstructing the central airway would provide a clinically meaningful improvement in net health outcome and whether the use is consistent with generally accepted medical practice. In response to requests, clinical input was received from 9 respondents, including 2 specialty society-level responses, 3 physician-level responses identified by specialty societies, and 4 physicians identified by 1 health system.

For individuals with NSCLC who are either poor surgical candidates or who required palliation for a lesion obstructing the central airway who receive cryoablation, clinical input supports this use provides a clinically meaningful improvement in net health outcome and indicates this use is consistent with generally accepted medical practice.

2009 Input

In response to requests, input was received from 2 physician specialty societies (5 reviews) and from 2 academic medical centers (3 reviews) while this policy was under review in 2009. There was strong support for the use of cryoablation in the treatment of select patients with renal tumors. There also was support for its use in the treatment of benign breast disease. Reviewers generally agreed cryoablation was investigational in the treatment of pancreatic cancer.

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American College of Radiology

The American College of Radiology Appropriateness Criteria (2009, updated 2021) for posttreatment follow-up and active surveillance of renal cell carcinoma [RCC] indicated that "Ablative therapies, such as radiofrequency ablation, microwave ablation, and cryoablation, have been shown to be effective and safe alternatives [to surgical resection] for the treatment of small, localized RCCs." These recommendations are based on a review of the data and expert consensus.

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisianais an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00023 Original Effective Date: 01/26/2004 Current Effective Date: 08/12/2024

American Urological Association

The American Urological Association (2021) updated its guidelines on the evaluation and management of clinically localized sporadic renal masses suspicious for renal cell carcinoma. The guideline statements on thermal ablation (radiofrequency ablation, cryoablation) are listed in Table 1.

Recommendations	LOR	LOE
Guideline statement 25		
Clinicians should consider thermal ablation (TA) as an alternate approach for the management of cT1a renal masses <3 cm in size. For patients who elect TA, a percutaneous technique is preferred over a surgical approach whenever feasible to minimize morbidity.	Moderate	С
Guideline statement 26		
Both radiofrequency ablation (RFA) and cryoablation may be offered as options for patients who elect thermal ablation.	Conditional	C
Guideline statement 28		
Counseling about thermal ablation should include information regarding an increased likelihood of tumor persistence or local recurrence after primary thermal ablation relative to surgical excision, which may be addressed with repeat ablation if further intervention is elected	Strong	В

Table 1. Guidelines on Localized Masses Suspicious for Renal Cell Carcinoma

LOE: level of evidence; LOR: level of recommendation.

National Comprehensive Cancer Network

Kidney Cancer

The National Comprehensive Network (NCCN) (v.4.2024) guidelines on kidney cancer state that "thermal ablation (eg, cryosurgery, radiofrequency ablation, microwave ablation) is an option for the management of clinical stage T1 renal lesions. Thermal ablation is an option for clinical T1b masses in select patients not eligible for surgery. Biopsy of lesions is recommended to be done prior

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisianais an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00023 Original Effective Date: 01/26/2004 Current Effective Date: 08/12/2024

to or at time of ablation. Ablative techniques may require multiple treatments to achieve the same local oncologic outcomes as conventional surgery."

The NCCN guidelines also note that "active surveillance and ablative techniques such as cryotherapy, microwave ablation, or radiofrequency ablation are alternative strategies for selected patients, particularly for those who are older, those with competing health risks, and those with T1b masses not eligible for surgery." Additionally, the guidelines note that "randomized phase III comparison of ablative techniques with surgical resection (ie, radical or partial nephrectomy by open or laparoscopic techniques) has not been performed."

Non-Small Cell Lung Cancer

The NCCN (6.2024) guidelines for NSCLC made the following relevant recommendations:

- Resection is the preferred local treatment modality for medically operable disease.
- Image-guided thermal ablation (IGTA) techniques include radiofrequency ablation, microwave ablation, and cryoablation.
- IGTA may be an option for select patients not receiving stereotactic ablative radiotherapy or definitive radiotherapy.
- IGTA may be considered for those patients who are deemed "high risk"- those with tumors that are for the most part surgically resectable but rendered medically inoperable due to comorbidities. In cases where IGTA is considered for high-risk or borderline operable patients, a multidisciplinary evaluation is recommended.
- IGTA is an option for the management of NSCLC lesions <3 cm. Ablation for NSCLC lesions >3 cm may be associated with higher rates of local recurrence and complications.
- The guidelines do not separate out recommendations by ablation technique and note that "each energy modality has advantages and disadvantages. Determination of energy modality to be used for ablation should take into consideration the size and location of the target tumor, risk of complication, as well as local expertise and/or operator familiarity."

Cancer Pain

The NCCN Guidelines on Adult Cancer Pain (v.2.2024) do not address cryoablation specifically for pain due to bone metastases, but note that "ablation techniques may...be helpful for pain management in patients who receive inadequate relief from pharmacological therapy."

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisianais an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00023 Original Effective Date: 01/26/2004 Current Effective Date: 08/12/2024

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in Table 2.

NCT No.	Trial Name	Planned Enrollment	Completion Date
Ongoing			
Renal cancer			
NCT02399124 ^a	ICESECRET PROSENSE ^{™‡} Cryotherapy for Renal Cell Carcinoma Trial	120	Feb 2026
NCT04506671	A Prospective, Non-randomized, in Parallel Groups Study Evaluating the Efficacy and Safety of Percutaneous Cryoablation and Partial Nephrectomy in Localized T1b Renal Tumor	142	Jun 2025
Breast cancer			
NCT05505643	COOL-IT: Cryoablation vs Lumpectomy in T1 Breast Cancers: A Randomized Controlled Trial With Safety Lead-in	256	Dec 2030
NCT04334785	Evaluation for the Effectiveness and Safety of Cryo-ablation in the Treatment of Early Invasive Breast Cancer	186	May 2025

Table 2. Summary of Key Trials

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisianais an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00023 Original Effective Date: 01/26/2004 Current Effective Date: 08/12/2024

Bone cancer			
NCT05615545	Safety and Efficacy of Cryoablation in the Treatment of Advanced Bone and Soft Tissue Tumors: a Single-center Retrospective Study	30	Oct 2023

NCT: national clinical trial.

^a Denotes industry-sponsored or cosponsored trial.

References

- 1. Yanagisawa T, Mori K, Kawada T, et al. Differential efficacy of ablation therapy versus partial nephrectomy between clinical T1a and T1b renal tumors: A systematic review and meta-analysis. Urol Oncol. Jul 2022; 40(7): 315-330. PMID 35562311
- 2. Uhlig J, Strauss A, Rücker G, et al. Partial nephrectomy versus ablative techniques for small renal masses: a systematic review and network meta-analysis. Eur Radiol. Mar 2019; 29(3): 1293-1307. PMID 30255245
- 3. Klatte T, Shariat SF, Remzi M. Systematic review and meta-analysis of perioperative and oncologic outcomes of laparoscopic cryoablation versus laparoscopic partial nephrectomy for the treatment of small renal tumors. J Urol. May 2014; 191(5): 1209-17. PMID 24231845
- 4. Tang K, Yao W, Li H, et al. Laparoscopic renal cryoablation versus laparoscopic partial nephrectomy for the treatment of small renal masses: a systematic review and meta-analysis of comparative studies. J Laparoendosc Adv Surg Tech A. Jun 2014; 24(6): 403-10. PMID 24914926
- 5. Andrews JR, Atwell T, Schmit G, et al. Oncologic Outcomes Following Partial Nephrectomy and Percutaneous Ablation for cT1 Renal Masses. Eur Urol. Aug 2019; 76(2): 244-251. PMID 31060824
- Rembeyo G, Correas JM, Jantzen R, et al. Percutaneous Ablation Versus Robotic Partial Nephrectomy in the Treatment of cT1b Renal Tumors: Oncologic and Functional Outcomes of a Propensity Score-weighted Analysis. Clin Genitourin Cancer. Apr 2020; 18(2): 138-147. PMID 31982346
- Yan S, Yang W, Zhu CM, et al. Comparison among cryoablation, radiofrequency ablation, and partial nephrectomy for renal cell carcinomas sized smaller than 2cm or sized 2-4cm: A population-based study. Medicine (Baltimore). May 2019; 98(21): e15610. PMID 31124938

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00023 Original Effective Date: 01/26/2004 Current Effective Date: 08/12/2024

- Pecoraro A, Palumbo C, Knipper S, et al. Cryoablation Predisposes to Higher Cancer Specific Mortality Relative to Partial Nephrectomy in Patients with Nonmetastatic pT1b Kidney Cancer. J Urol. Dec 2019; 202(6): 1120-1126. PMID 31347950
- 9. Cronan J, Dariushnia S, Bercu Z, et al. Systematic Review of Contemporary Evidence for the Management of T1 Renal Cell Carcinoma: What IRs Need to Know for Kidney Cancer Tumor Boards. Semin Intervent Radiol. Aug 2019; 36(3): 194-202. PMID 31435127
- Morkos J, Porosnicu Rodriguez KA, Zhou A, et al. Percutaneous Cryoablation for Stage 1 Renal Cell Carcinoma: Outcomes from a 10-year Prospective Study and Comparison with Matched Cohorts from the National Cancer Database. Radiology. Aug 2020; 296(2): 452-459. PMID 32515677
- 11. Stacul F, Sachs C, Giudici F, et al. Cryoablation of renal tumors: long-term follow-up from a multicenter experience. Abdom Radiol (NY). Sep 2021; 46(9): 4476-4488. PMID 33912986
- 12. Lee SH, Choi WJ, Sung SW, et al. Endoscopic cryotherapy of lung and bronchial tumors: a systematic review. Korean J Intern Med. Jun 2011; 26(2): 137-44. PMID 21716589
- 13. Niu L, Xu K, Mu F. Cryosurgery for lung cancer. J Thorac Dis. Aug 2012; 4(4): 408-19. PMID 22934144
- Callstrom MR, Woodrum DA, Nichols FC, et al. Multicenter Study of Metastatic Lung Tumors Targeted by Interventional Cryoablation Evaluation (SOLSTICE). J Thorac Oncol. Jul 2020; 15(7): 1200-1209. PMID 32151777
- 15. de Baere T, Tselikas L, Woodrum D, et al. Evaluating Cryoablation of Metastatic Lung Tumors in Patients--Safety and Efficacy: The ECLIPSE Trial--Interim Analysis at 1 Year. J Thorac Oncol. Oct 2015; 10(10): 1468-74. PMID 26230972
- 16. Moore W, Talati R, Bhattacharji P, et al. Five-year survival after cryoablation of stage I nonsmall cell lung cancer in medically inoperable patients. J Vasc Interv Radiol. Mar 2015; 26(3): 312-9. PMID 25735518
- 17. Ratko TA, Vats V, Brock J, et al. Local Nonsurgical Therapies for Stage I and Symptomatic Obstructive Non-Small-Cell Lung Cancer (AHRQ Comparative Effectiveness Review No. 112). Rockville, MD: Agency for Healthcare Research and Quality; 2013.
- 18. Maiwand MO, Asimakopoulos G. Cryosurgery for lung cancer: clinical results and technical aspects. Technol Cancer Res Treat. Apr 2004; 3(2): 143-50. PMID 15059020
- 19. Zhao Z, Wu F. Minimally-invasive thermal ablation of early-stage breast cancer: a systemic review. Eur J Surg Oncol. Dec 2010; 36(12): 1149-55. PMID 20889281

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisianais an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00023 Original Effective Date: 01/26/2004 Current Effective Date: 08/12/2024

- 20. Simmons RM, Ballman KV, Cox C, et al. A Phase II Trial Exploring the Success of Cryoablation Therapy in the Treatment of Invasive Breast Carcinoma: Results from ACOSOG (Alliance) Z1072. Ann Surg Oncol. Aug 2016; 23(8): 2438-45. PMID 27221361
- 21. Niu L, Mu F, Zhang C, et al. Cryotherapy protocols for metastatic breast cancer after failure of radical surgery. Cryobiology. Aug 2013; 67(1): 17-22. PMID 23619024
- 22. Manenti G, Perretta T, Gaspari E, et al. Percutaneous local ablation of unifocal subclinical breast cancer: clinical experience and preliminary results of cryotherapy. Eur Radiol. Nov 2011; 21(11): 2344-53. PMID 21681574
- 23. Pusztaszeri M, Vlastos G, Kinkel K, et al. Histopathological study of breast cancer and normal breast tissue after magnetic resonance-guided cryotherapy ablation. Cryobiology. Aug 2007; 55(1): 44-51. PMID 17604016
- 24. Sabel MS, Kaufman CS, Whitworth P, et al. Cryoablation of early-stage breast cancer: work-inprogress report of a multi-institutional trial. Ann Surg Oncol. May 2004; 11(5): 542-9. PMID 15123465
- 25. Tanaka S. Cryosurgical treatment of advanced breast cancer. Skin Cancer. Jan 1995;10:9-18.
- 26. Pfleiderer SO, Freesmeyer MG, Marx C, et al. Cryotherapy of breast cancer under ultrasound guidance: initial results and limitations. Eur Radiol. Dec 2002; 12(12): 3009-14. PMID 12439583
- 27. Suzuki Y. Cryosurgical treatment of advanced breast cancer and cryoimmunological responses. Skin Cancer. 1995;10:19-26.
- 28. Morin J, Traoré A, Dionne G, et al. Magnetic resonance-guided percutaneous cryosurgery of breast carcinoma: technique and early clinical results. Can J Surg. Oct 2004; 47(5): 347-51. PMID 15540687
- 29. Kaufman CS, Bachman B, Littrup PJ, et al. Office-based ultrasound-guided cryoablation of breast fibroadenomas. Am J Surg. Nov 2002; 184(5): 394-400. PMID 12433600
- 30. Kaufman CS, Littrup PJ, Freman-Gibb LA, et al. Office-based cryoablation of breast fibroadenomas: 12-month followup. J Am Coll Surg. Jun 2004; 198(6): 914-23. PMID 15194073
- 31. Kaufman CS, Bachman B, Littrup PJ, et al. Cryoablation treatment of benign breast lesions with 12-month follow-up. Am J Surg. Oct 2004; 188(4): 340-8. PMID 15474424
- 32. Littrup PJ, Freeman-Gibb L, Andea A, et al. Cryotherapy for breast fibroadenomas. Radiology. Jan 2005; 234(1): 63-72. PMID 15550369
- 33. Kaufman CS, Littrup PJ, Freeman-Gibb LA, et al. Office-based cryoablation of breast fibroadenomas with long-term follow-up. Breast J. 2005; 11(5): 344-50. PMID 16174156

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisianais an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Policy # 00023 Original Effective Date: 01/26/2004 Current Effective Date: 08/12/2024

- 34. Nurko J, Mabry CD, Whitworth P, et al. Interim results from the FibroAdenoma Cryoablation Treatment Registry. Am J Surg. Oct 2005; 190(4): 647-51; discussion 651-2. PMID 16164941
- 35. Tao Z, Tang Y, Li B, et al. Safety and effectiveness of cryosurgery on advanced pancreatic cancer: a systematic review. Pancreas. Jul 2012; 41(5): 809-11. PMID 22695092
- 36. Keane MG, Bramis K, Pereira SP, et al. Systematic review of novel ablative methods in locally advanced pancreatic cancer. World J Gastroenterol. Mar 07 2014; 20(9): 2267-78. PMID 24605026
- 37. Li J, Chen X, Yang H, et al. Tumour cryoablation combined with palliative bypass surgery in the treatment of unresectable pancreatic cancer: a retrospective study of 142 patients. Postgrad Med J. Feb 2011; 87(1024): 89-95. PMID 21131612
- Xu KC, Niu LZ, Hu YZ, et al. A pilot study on combination of cryosurgery and (125)iodine seed implantation for treatment of locally advanced pancreatic cancer. World J Gastroenterol. Mar 14 2008; 14(10): 1603-11. PMID 18330956
- 39. Kovach SJ, Hendrickson RJ, Cappadona CR, et al. Cryoablation of unresectable pancreatic cancer. Surgery. Apr 2002; 131(4): 463-4. PMID 11935137
- 40. Meller I, Weinbroum A, Bickels J, et al. Fifteen years of bone tumor cryosurgery: a single-center experience of 440 procedures and long-term follow-up. Eur J Surg Oncol. Aug 2008; 34(8): 921-927. PMID 18158228
- 41. Callstrom MR, Dupuy DE, Solomon SB, et al. Percutaneous image-guided cryoablation of painful metastases involving bone: multicenter trial. Cancer. Mar 01 2013; 119(5): 1033-41. PMID 23065947
- 42. Jennings JW, Prologo JD, Garnon J, et al. Cryoablation for Palliation of Painful Bone Metastases: The MOTION Multicenter Study. Radiol Imaging Cancer. Mar 2021; 3(2): e200101. PMID 33817650
- 43. Purysko AS, Nikolaidis P, Khatri G, et al. ACR Appropriateness Criteria® Post-Treatment Follow-up and Active Surveillance of Clinically Localized Renal Cell Carcinoma: 2021 Update. J Am Coll Radiol. May 2022; 19(5S): S156-S174. PMID 35550799
- 44. American College of Radiology (ACR). ACR Appropriateness Criteria: Post-treatment followup and active surveillance of clinically localized renal cell carcinoma. Updated 2021. https://acsearch.acr.org/docs/69365/Narrative/.
- 45. Campbell SC, Clark PE, Chang SS, et al. Renal Mass and Localized Renal Cancer: Evaluation, Management, and Follow-Up: AUA Guideline: Part I. J Urol. Aug 2021; 206(2): 199-208. PMID 34115547

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisianais an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00023 Original Effective Date: 01/26/2004 Current Effective Date: 08/12/2024

46. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Kidney Cancer. Version 4.2024.

http://www.nccn.org/professionals/physician_gls/pdf/kidney.pdf.

- 47. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Non-Small Cell Lung Cancer. Version 6.2024. http://www.nccn.org/professionals/physician_gls/pdf/nscl.pdf.
- 48. National Comprehensive Cancer Network. Adult Cancer Pain. Version 2.2024. https://www.nccn.org/professionals/physician_gls/pdf/pain.pdf.

Policy History

I oney III.	
Original Effect	ive Date: 01/26/2004
Current Effecti	ve Date: 08/12/2024
10/21/2003	Medical Policy Committee review
01/26/2004	Managed Care Advisory Committee approval
12/07/2005	Medical Director review
12/20/2005	Medical Policy Committee review. Format revision. FDA approval information
	added to policy.
02/23/2006	Quality Care Advisory Council approval
10/10/2007	Medical Director review
10/17/2007	Medical Policy Committee approval. No change to coverage eligibility.
03/04/2009	Medical Director review
03/18/2009	Medical Policy Committee approval. Changed localized renal cell carcinoma from
	investigational to eligible for coverage with criteria. Breast fibroadenomas removed
	from this policy and made into a separate policy.
03/05/2010	Medical Policy Committee review
03/19/2010	Medical Policy Implementation Committee approval. Added benign tumors of the
	breast to be investigational.
03/03/2011	Medical Policy Committee review
03/16/2011	Medical Policy Implementation Committee approval. Renal cell carcinomas in
	patients who are surgical candidates was added as investigational.
03/01/2012	Medical Policy Committee review
03/21/2012	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisianais an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00023 Original Effective Date: 01/26/2004 Current Effective Date: 08/12/2024

- 03/07/2013 Medical Policy Committee review
- 03/20/2013 Medical Policy Implementation Committee approval. Title changed from "Cryosurgery Ablation of Miscellaneous Solid Tumors other than Liver or Prostate" to "Cryosurgery Ablation of Miscellaneous Solid Tumors Other than Liver or Prostate Tumors or Breast Fibroadenomas". Removed the second criteria bullet for treatment of renal cell carcinoma requiring that the patient not be considered as a surgical candidate due to co-morbid disease. Lung cancer added to investigational statement. The investigational statement was revised for clarification.
- 03/06/2014 Medical Policy Committee review
- 03/19/2014 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
- 08/03/2015 Coding update: ICD10 Diagnosis code section added; ICD9 Procedure code section removed.
- 09/03/2015 Medical Policy Committee review
- 09/23/2015 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
- 09/08/2016 Medical Policy Committee review
- 09/21/2016 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
- 01/01/2017 Coding update: Removing ICD-9 Diagnosis Codes
- 12/07/2017 Medical Policy Committee review
- 12/20/2017 Medical Policy Implementation Committee approval. Added a criteria bullet to localized renal cell carcinoma that states "the patient is not considered a surgical candidate." Added that cryosurgical ablation to treat lung cancer may be considered eligible for coverage with criteria. Added an investigational statement for lung cancer when criteria are not met. Replaced the investigational statement regarding cryosurgical ablation for malignant tumors with an investigational statement indicating that cryosurgical ablation as a treatment for benign or malignant tumors of the breast, pancreas, or bone and other solid tumors or metastases outside the liver, prostate and breast fibroadenomas is considered to be investigational.
- 01/01/2018 Coding update
- 08/31/2018 Coding update
- 12/06/2018 Medical Policy Committee review

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisianais an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00023 Original Effective Date: 01/26/2004 Current Effective Date: 08/12/2024 12/19/2018 Medical Policy Implementation Committee approval. Coverage eligibility unchanged. Medical Policy Committee review 12/05/2019 Medical Policy Implementation Committee approval. Coverage eligibility 12/11/2019 unchanged. Medical Policy Committee review 05/07/2020 05/13/2020 Medical Policy Implementation Committee approval. Coverage eligibility unchanged. Medical Policy Committee review 05/06/2021 Medical Policy Implementation Committee approval. Coverage eligibility 05/12/2021 unchanged. Medical Policy Committee review 07/01/2021 Medical Policy Implementation Committee approval. Title changed from 07/14/2021 "Cryosurgery Ablation of Miscellaneous Solid Tumors other than Liver or Prostate Tumors or Breast Fibroadenomas" to "Cryosurgical Ablation of Miscellaneous Solid Tumors other than Liver, Prostate, or Dermatologic Tumors". Revised the When Services May be Eligible and When Services Are Investigational sections. 07/07/2022 Medical Policy Committee review 07/13/2022 Medical Policy Implementation Committee approval. Title changed from "Cryosurgery Ablation of Miscellaneous Solid Tumors other than Liver, Prostate, or Dermatologic Tumors" to "Cryoablation of Tumors Located in the Kidney, Lung, Breast, Pancreas, or Bone.". "... and other solid tumors or metastases outside the liver and prostate" was removed from the investigational statement. Added Policy Guidelines. Medical Policy Committee review 03/02/2023 03/08/2023 Medical Policy Implementation Committee approval. Replaced "patients" with "individuals" in the coverage section. Added a suspected renal mass to be eligible for coverage with criteria for localized renal cell carcinoma. Changed a criteria bullet for cryosurgical ablation of a renal mass suspected to be localized renal cell carcinoma from "not considered a surgical candidate" to "considered a high-risk surgical candidate". 07/06/2023 Medical Policy Committee review

07/12/2023 Medical Policy Implementation Committee approval. Replaced renal mass section with "Based on review of available data, the Company may consider cryosurgical

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisianais an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00023 Original Effective Date: 01/26/2004 Current Effective Date: 08/12/2024

ablation for clinically localized, suspected renal malignancy for individuals with peripheral lesions that are less than or equal to 4 cm in diameter to be eligible for coverage."

- 07/02/2024 Medical Policy Committee review
- 07/10/2024 Medical Policy Implementation Committee approval. Moved coverage for renal malignancy to a "When Services Are Eligible for Coverage" section for clarity. Revised the investigational statement for cryosurgical ablation to treat localized renal cell carcinoma for clarity. Updated NCCN Guidelines to Version 4.2024 for Kidney Cancer in the Supplemental Information section. Coverage eligibility unchanged.

Next Scheduled Review Date: 07/2025

Coding

The five character codes included in the Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology $(CPT^{\circledast})^{\ddagger}$, copyright 2023 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.

The responsibility for the content of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines is with Blue Cross and Blue Shield of Louisiana and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.

CPT is a registered trademark of the American Medical Association.

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisianais an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00023 Original Effective Date: 01/26/2004 Current Effective Date: 08/12/2024

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	0581T, 19105, 20983, 32994, 50250, 50542, 50593
HCPCS	C2618
ICD-10 Diagnosis	All related diagnoses

*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
 - 1. Consultation with technology evaluation center(s);
 - 2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
 - 3. Reference to federal regulations.

**Medically Necessary (or "Medical Necessity") - Health care services, treatment, procedures, equipment, drugs, devices, items or supplies that a Provider, exercising prudent clinical judgment, would provide to a patient for the purpose of preventing, evaluating, diagnosing or treating an illness, injury, disease or its symptoms, and that are:

- A. In accordance with nationally accepted standards of medical practice;
- B. Clinically appropriate, in terms of type, frequency, extent, level of care, site and duration, and considered effective for the patient's illness, injury or disease; and

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00023 Original Effective Date: 01/26/2004 Current Effective Date: 08/12/2024

C. Not primarily for the personal comfort or convenience of the patient, physician or other health care provider, and not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of that patient's illness, injury or disease.

For these purposes, "nationally accepted standards of medical practice" means standards that are based on credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community, Physician Specialty Society recommendations and the views of Physicians practicing in relevant clinical areas and any other relevant factors.

‡ Indicated trademarks are the registered trademarks of their respective owners.

NOTICE: If the Patient's health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

NOTICE: Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage.

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisianais an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.